AD-A189 228 LARGE MOMENTUM PAIRING IN OME-DIMENSIONAL SYSTEMS(U)


LENY STATE UNIV ONTO LIQUID CRYSTAL INST BY ALLEMDER

UMCLASSIFIED

LARGE MOMENTUM PAIRING IN OME-DIMENSIONAL SYSTEMS(U)

LENY STATE UNIV ONTO LIQUID CRYSTAL INST BY ALLEMDER

F/G 20/5 NL



•

AD A	EPORT DOCUM	MENTATION		-		
AD-A189	228	16. RESTRICTIVE	MARKINGS	G FILE	COPY	
2a. SECURITY CLASSIFICATION ACTION.		AVAILABILITY OF				
2b. DECLASSIFICATION / DOWNGRADING	Approved for public release, distribution unlimited					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)				
		AFO	SR-TR- 8	37-17	24	
6a. NAME OF PERFORMING ORGANIZATI	ON 6b. OFFICE SYMBOL (If applicable)		NITORING ORGA			
Kent State University	(ii appiicaule)	AFOSR				
6c. ADDRESS (City, State, and ZIP Code)	7b ADDRESS (City, State, and ZIP Code)					
Department of Physics as Kent State University	Bidg 410					
Kent , Ohio 44242	Bolling AFB, DC 20332-6448					
8a. NAME OF FUNDING/SPONSORING 8b. OFFICE SYMBO		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER				
ORGANIZATION (If applicable) AFOSR NP		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-84-0185 10 SOURCE OF FUNDING NUMBERS				
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF F		- 62	VW A	
		PROGRAM ELEMENT NO	PROJECT NO	TASK NO.	WORK UNIT ACCESSION NO.	
same as 7b.		61102F	2301	A8 💝	COE	
11. TITLE (Include Security Classification)						
"LARGE MOMENTUM PAIRING	IN ONE-DIMENSIONAL S	YSTEMS" (U)		·		
12 PERSONAL AUTHOR(S) Dr. D. W. Allender						
13a. TYPE OF REPORT 13b. FINAL FR	14. DATE OF REPO	RT (Year Month, L	Pay) 15. PAGE	COUNT 4		
16. SUPPLEMENTARY NOTATION						
	<u></u>					
17. COSATI CODES	the state of the s					
FIELD GROUP SUB-GRO		NTUM, PAIRIN	G			
BAROL, HOLDHIOH, LAIRING						
19. ABSTRACT (Continue on reverse if necessary and identify by block number)						
The first and primary objective was to determine the correlation function describing large momentum pairing. A comparison of this correlation function with other corre-						
lations (such as singlet pairing, triplet pairing, charge density wave, and spin density						
wave) is now pCossible.) The second objective was to strengthen and maintain a						
collaboration with researchers D.C. Reynolds and S.B. Nam at Wright-Patterson AFB.						
In light of the recent discovery of high temperature superconductivity (T(C) 95K) in rare earth copper oxide materials, it is apparent that a revolution in superconductivity						
research is beginning. This area will be of great importance to the Air Force in the						
future; thus close ties	S between University	researchers	and Air Ford	e laborato	ries will	
be practical and benef	ICIAL.					
	N. Carlotte					
20. DISTRIBUTION / AVAILABILITY OF AB	21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED					
☑ UNCLASSIFIED/UNLIMITED ☑ SA 22a. NAME OF RESPONSIBLE INDIVIDUAL	UNCLASSIFIED 22b. TELEPHONE (Include Area Code) 22c. OFFICE SYMBOL					
DR. ROBERT J. BARKER (202)767-5011				NP		
DD FORM 1473, 84 MAR	83 APR edition may be used un		SECURITY	CLASSIFICATION	OF THIS PAGE	
	All other editions are of	osolete.		OTETED		

By 20 20 1 UNCLASSIFIED

AFOSR Grant Number 84-0185 Final Technical Report September 1987

AFOSR-TR- 87-1792

LARGE MOMENTUM PAIRING IN ONE-DIMENSIONAL SYSTEMS

Dr. D.W. Allender
Department of Physics and Liquid Crystal Institute
Kent State University
Kent, Ohio 44242

[Telephone: (216) 672-2577]

Access	ion Fo	r		
NTIS	GRA&I			
DTIC 1	CAB	1		
Unannounced				
Justification				
By				
1 1	Avail and/or			
Dist	Special			
A-1				

Controlling Office: USAF Office of Scientific Research/NP Bolling Air Force Base, DC 20332



1. Introduction and Statement of Accomplishments

This report summarizes the progress that was made on a theoretical investigation of large momentum pairing in one-dimensional superconductors that was proposed and supported by grant AFOSR-84-0185. The grant period from July 1984 to December 1985 was productive and successful in that two major objectives of the proposed research were achieved.

The first and primary objective was to determine the correlation function describing large momentum pairing. A comparison of this correlation function with other correlations (such as singlet pairing, triplet pairing, charge density wave, and spin density wave) is now possible. The second objective was to strengthen and maintain a collaboration with researchers D.C. Reynolds and S.B. Nam at Wright-Patterson AFB. In light of the recent discovery of high temperature superconductivity ($T_C \sim 95 K$) in rare earth copper oxide materials, it is apparent that a revolution in superconductivity research is beginning. This area will be of great importance to the Air Force in the future; thus close ties between University researchers and Air Force laboratories will be practical and beneficial.

In the remainder of this report, details are given about the first objective, and the facts regarding activities supported by this grant are presented.

II. The Large Momentum Pairing Correlation

In a review article in Highly Conducting One-Dimensional Solids (Plenum Press, NY, 1979), V.J. Emery presented a clear discussion of the theory of the one-dimensional electron gas. The model Hamiltonian he considered is often called the extended Hubbard model, and consists of three parts: (1) H_0 describing hopping from site to site, (2) H_1 giving on-site interaction, and (3) H_2 presenting intersite

interaction. H_1 and H_2 are mostly due to Coulomb effects, but phonons and other phenomena also contribute.

The standard correlation functions were described by Emery, and can be written as

$$\chi_{\lambda}^{R} = -i \theta(t) \langle \left[\hat{O}_{\lambda}^{+}(xt), \hat{O}_{\lambda}(00) \right] \rangle$$

where $\lambda = SP$ (singlet pairing), TP (triplet pairing), CDW (charge density wave), and SDW (spin density wave). The \hat{O}_{λ} operators are

$$\hat{O}_{SP} = \psi_{2\sigma}^+ \psi_{1-\sigma}^+$$

$$\dot{O}_{TP} = \psi_{2\sigma}^+ \psi_{1\sigma}^+$$

$$\dot{O}_{CDW} = \frac{1}{\sqrt{2}} \left[\psi_{2\uparrow}^+ \psi_{1\uparrow}^- + \psi_{2\downarrow}^+ \psi_{1\downarrow}^- \right]$$

$$\hat{O}_{SDW} = \psi_{2\sigma}^+ \psi_{1-\sigma}$$

The $\psi_{i\sigma}$ operators are the usual second-quantized fermion field operators. Label i=1 implies a particle with momentum to the right, label i=2 means momentum to the left, and σ denotes spin. The results found by evaluating these correlation functions using the full model Hamiltonian and the boson representation technique were

$$x_{\lambda}^{R} = \frac{\theta(t)}{2\sigma^{2}r^{2}} Im \left\{ \left[r^{2}K\left(x - v_{s}^{\prime}t, r\right)K^{*}\left(x + v_{s}^{\prime}t, r\right) \right]^{(\theta_{s}^{\pm 1})/2} \right\}$$

$$\times \left[r^2 K \left(x - v \cdot t, r \right) K \left(x + v \cdot t, r \right) \right]^{(\theta_c^{\pm 1})/2} \right\}.$$

The notation and functions in these equations are too cumbersome to carefully define in this report, but they are given by Emery. The important thing to notice is that the exponents $\theta_s^{\pm 1}$ and $\theta_c^{\pm 1}$ are functions of the parameters of the Hamiltonian.

The theoretical work carried out under this grant was a calculation of a new and previously unrecognized correlation function. Using the notation $\lambda = LMP$ (for large momentum pairing), and the definition

$$\hat{O}_{LMP} = \psi_{1\uparrow}^+ \psi_{1\downarrow}^+$$

the correlation function χ^R_{LMP} was evaluated by means of the boson representation. The answer is

$$\chi_{LMP} = \frac{\theta(t)}{2\pi^2} Im \left\{ K^2 \left(x - v_c^{\dagger} t, \alpha \right) \left[r^2 K \left(x - v_c^{\dagger} t, r \right) K^* \left(x + v_c^{\dagger} t, r \right) \right]^{2\gamma} \right\}$$

where

$$2\gamma = -1 + \left(1 - \frac{\overline{W}_{\parallel}^2}{4}\right)^{-1/2},$$

and all other notation is given by Emery (\bar{W}_i is a parameter of the Hamiltonian).

The significance of this expression is that, by Fourier transforming, the exponent of the frequency dependence can be found. If the exponent is negative, the correlation diverges and one expects the ground state to have large momentum pairs formed. The Hamiltonian parameter values for which the exponent is negative can in principle be mapped out. These calculations to find the exponent have <u>not</u> been done, but are now possible. Further research effort to carry out these calculations is necessary.

III. Publications

The results of this research have not been published to date, although a manuscript is in preparation which presents the derivation of the results stated above in section II.

IV. Personnel

The research described here was pursued at Kent State University by the principal investigator, D.W. Allender. Two graduate students, Kazi Motakabbir and Young Hee Lee also carried out research and received support. Kazi received his Ph.D. in December 1985 and Young Hee completed his in August 1985.

V. Interactions

The primary interaction supported by this grant was the collaboration between D.C. Reynolds and S.B. Nam at Wright-Patterson AFB and the principal investigator. One trip was made to Wright-Patterson AFB where useful discussions were held.

Also supported was a presentation at the Midwest Solid State Theory Symposium in Minneapolis, MN in September 1984, as well as participation in two other conferences: the March 1985 meeting of the American Physical Society in Baltimore, MD and the Materials and Mechanisms of Superconductivity Conference in Ames, IA in May 1985.

DATED AND STREET OF THE PROPERTY OF THE PROPER